

## In-Medium strangeness dynamics at $\bar{\text{PANDA}}^*$

T. Gaitanos<sup>1</sup> and H. Lenske<sup>1,2</sup>

<sup>1</sup>Inst. für Theor. Physik, Universität Giessen, Germany; <sup>2</sup>GSI Darmstadt, Germany

Hypernuclear physics opens a unique opportunity to explore the hyperon-nucleon ( $YN$ ) and the hyperon-hyperon ( $YY$ ) in-medium interactions at terrestrial laboratories. Such studies are important for a better understanding of the in-medium interactions with strangeness degree of freedom, which is still uncertain, but of relevance for nuclear astrophysics [1]. The experimental knowledge on multi-strangeness hypernuclei has been so far scarce. However, recent experiments at GSI are very promising. In the Hy-phi [2] experiment precise production rates of single- $\Lambda$  ( $S = -1$ ) hypernuclei were measured. An abundant production of double- $\Lambda$  ( $S = -2$ ) clusters is also expected in the  $\bar{\text{PANDA}}$  experiment [3] at FAIR.

We investigate theoretically the multi-strangeness dynamics within the Giessen-BUU (GiBUU) approach [4]. In particular, we have studied antiproton-induced reactions allowing for reactions of the secondary  $\Xi$ -beam on a second target. The formation of double- $\Lambda$  hypernuclei occurs in the  $\Xi$ -interaction with the second target [5]. Two issues are of importance for the production of double-strangeness hypermatter, see Fig. 1. At first, the absorption of the  $\Xi$ -beam inside the target matter decreases with increasing energy. This is due to the strong decrease of the elementary  $\Xi N \rightarrow \Lambda\Lambda$  channel [5]. Secondly, the abundance of bound  $\Lambda$ -hyperons also strongly decreases with rising incident  $\Xi$ -energy. This is mainly due to the repulsive vector field which becomes more pronounced as the particle energy increases. These effects lead to a rather strong energy dependent rise of the double- $\Lambda$  hypernuclear production cross section. Thus low-energetic cascade beams should be used at  $\bar{\text{PANDA}}$  in order to obtain high production rates of double-strangeness hyperfragments [5].

So far bare interactions for elementary  $YN, YY$ -channels have been used in transport approaches. However, at  $\bar{\text{PANDA}}$  the strangeness dynamics takes place at densities closely up to saturation. We have studied in-medium effects on various  $YN$ -processes by solving the Lippmann-Schwinger equation with a Pauli-exclusion operator for intermediate states as the leading-order in-medium effect [6, 7]. The elementary  $YN$  cross sections are indeed influenced by in-medium effects, in particular, at low energies, as shown in Fig. 2 (similar effects occur for quasi-elastic channels with strangeness exchange).

In the  $S=-2$  sector the situation is still very controversial theoretically [8]. Presently we are investigating the influence of various calculations on  $S=-2$ -channels [9] on the strangeness production in  $\bar{p}$ -induced reactions and extending our studies to the  $S=-3$  sector by accounting for the




Figure 1: GiBUU results for the time dependence of all and free  $\Lambda$ s and of  $\Xi$ s for central  $\Xi$ -induced reactions on Cu-target at beam energies as indicated [5].

formation of  $\Omega$ -baryons [10]. The preliminary results are very promising concerning a possible formation of multi-strangeness bound systems at  $\bar{\text{PANDA}}$ . We emphasize the relevance of our theoretical results for the future activities at FAIR.




Figure 2: Energy dependence of elastic in-medium cross sections for  $\Lambda N$  and  $\Sigma N$  scattering at various densities as indicated [7].

## References

- [1] J. Schaffner, I. Mishustin, Phys. Rev. C53 (1996) 1416.
- [2] C. Rappold, *et al.*, Nucl. Phys. A622 (2010) 231.
- [3] A.S. Lorente, *et al.*, Phys. Lett. B 697 (2011) 222.
- [4] O. Buss, *et al.*, Phys. Rep. 512 (2012) 1.
- [5] T. Gaitanos, *et al.*, Nucl. Phys. A881 (2012) 240.
- [6] T. Gaitanos, *et al.*, Nucl. Phys. A914 (2013) 405.
- [7] A. Obermann, Masters Thesis, 2011, Univ. Giessen, unpublished.
- [8] J.K. Ahn, *et al.*, Phys. Lett. B 633 (2006) 214.
- [9] T.A. Rijken, Y. Yamamoto, nucl-th/0608074; Y. Fujiwara, *et al.*, Phys. Rev. C64 (2001) 054001.
- [10] T. Gaitanos, *et al.*, in preparation.

\* Work supported by BMBF, DFG, HIC for FAIR, and GSI-JLU Giessen collaboration agreement